


## Feeding recommendations



These recommendations are given for indicative purposes. They correspond to the needs of the genetics to make it express all its potential.

The quality of the feed will be essentially linked to the raw materials used in its formulation.


### ➤ Program with 3 feeds



*Number of days before (-) or after (+) kindling*

	LACTATION	AROUND WEANING	FINISHING
<b>Digestible Energy (kcal/kg)</b>	2600 - 2650	2350 - 2400	2450 - 2500
<b>Gross Protein (%)</b>	17,5 - 18	15,5 - 16	15 - 15,5
<b>Fat (%)</b>	3,2 - 3,5	3 - 3,2	3 - 3,2
<b>Gross Fiber (%)</b>	14,5 - 15	17 - 17,5	15,5 - 16
<b>Starch (%)</b>	15 - 16	7 - 8	11 - 12
<b>Lignine (%)</b>	4,5	6,5 - 7	5,5 - 6
<b>Lysine (%)</b>	0,85	0,7	0,7
<b>Methionine + Cystine (%)</b>	0,7	0,6	0,6

### ➤ Program with 4 feeds



*Number of days before (-) or after (+) kindling*

	LACTATION	PRE WEANING	GROWING	FINISHING
<b>Digestible Energy (kcal/kg)</b>	2650 - 2700	2500 - 2550	2350 - 2400	2450 - 2500
<b>Gross Protein (%)</b>	17,5 - 18	16,8 - 17,2	15,5 - 16	15 - 15,5
<b>Fat (%)</b>	3,2 - 3,5	3 - 3,2	3 - 3,2	3 - 3,2
<b>Gross Fiber (%)</b>	14 - 15	15 - 16	17 - 17,5	15,5 - 16
<b>Starch (%)</b>	16 - 17	11 - 12	7 - 8	11 - 12
<b>Lignine (%)</b>	4,5	5 - 5,5	6,5 - 7	5,5 - 6
<b>Lysine (%)</b>	0,85	0,72	0,7	0,7
<b>Methionine + Cystine (%)</b>	0,7	0,65	0,6	0,6



The features of the Pre Weaning feed are those that fit the best with the preparation program of young females.



For the females at 1<sup>st</sup> birth, do not use the lactation feed of the program with 4 feeds.